

## ARTICLE

# Measurement properties of the Illness Perception Questionnaire in practitioners working with patients with co-morbid schizophrenia and learning disability

Colin Martin BSc MSc PhD MBA RN YCAP FHEA CPsychol CSci AFBPsS<sup>a</sup>, Mick Fleming RN BA MA PhD YCAP<sup>b</sup> and Stewart McCorkindale RN MSc<sup>c</sup>

a Professor of Mental Health, Faculty of Society and Health, Buckinghamshire New University, Uxbridge Campus, Middlesex, UK

b Senior Lecturer in Mental Health, University of the West of Scotland, Ayr, Scotland, UK

c Team Leader, Inverclyde Community Learning Disability Team, Greenock, Scotland, UK

## Abstract

**Rationale, aims and objectives:** The widely-used Illness Perception Questionnaire (IPQ) adapted for use in patients with major mental health problems and those who care and work with them, was originally developed for evaluating the illness perceptions of patients with physical pathology. There have been concerns expressed regarding the appropriateness of the IPQ and derivative measures of it in axis I disorders such as schizophrenia. The current investigation examined the measurement characteristics of a modified version of the IPQ in mental health practitioners working with patients with schizophrenia co-morbid with learning disability.

**Method:** Two hundred and ten mental health practitioners working with patients with schizophrenia co-morbid with learning disability participated in the study. Contemporary models of the measurement structure of the IPQ were evaluated using confirmatory factor analysis. The internal reliability of the IPQ was also evaluated.

**Results:** Single factor, 5-factor and 6-factor models were evaluated against data. All models were found to offer a poor fit to the data. The internal reliability of the measure was also found to be unsatisfactory overall.

**Conclusions:** Consistent with one previous study that has found poor measurement characteristics of the IPQ when used in the context of schizophrenia, the current study found the measure to be a poorly performing index of illness perceptions. Echoing previous observations of the use of this tool in axis I disorder, it is recommended that a new measure of illness perceptions be developed using contemporary research methodology sensitive to the context of psychotic illness in order to provide a more effective tool for the provision of person-centered psychiatry and healthcare.

## Keywords

Evaluation, Illness Perception Questionnaire (IPQ), learning disabilities/difficulties, measurement properties, methodological robustness, person-centered healthcare, person-centered psychiatry reliability, psychometric properties, schizophrenia, validity

## Correspondence address

Professor Colin R. Martin, Faculty of Society and Health, Buckinghamshire New University, Uxbridge Campus, 106 Oxford Road, Uxbridge, Middlesex, UB8 1NA, UK. E-mail: colin.martin@bucks.ac.uk

Accepted for publication: 7 September 2015

## Introduction

It is increasingly recognized, within the context of the provision of comprehensive patient care, that the role of illness perceptions are of critical importance to the course and outcomes of a range of pathologies [1-5]. Appreciation of the role of illness perceptions to outcome has not only enlightened the opportunities for holistic, person-centered care delivery [4,6-9], but has also precipitated awareness of the potential for interventions that attempt to positively influence clinical outcomes by improving illness perceptions directly [1,10-13]. Mechanisms that mediate the influence and role of illness perceptions on clinical

outcome remain largely theoretical and to a degree speculative [14-18]. However, the evidence from a range of clinical studies highlights, incontrovertibly, a clinically pertinent and relevant role for these mechanisms. One of the main and certainly most widely used measures of illness perceptions is the Illness Perception Questionnaire [5]. The IPQ was developed within the theoretical context of Leventhal's self-regulatory model to be a conceptually and psychometrically accurate multi-dimensional measure of self-report patient illness appraisal. Five illness perception domains were envisaged to comprise the main illness perception factors of relevance to patients, these

being: (i) illness identity; (ii) cause; (iii) timeline; (iv) consequences & (v) control/cure.

Thirty-six items were included in the original IPQ [5]. Initial evaluation of the basic measurement characteristics of the IPQ was overall positive and promising, but, recognising some less than optimal measurement properties, the instrument developers suggested further work to evaluate the tool [5]. Evaluation of the measurement characteristics of the IPQ has been extensive and led directly to the development of a revised version of the tool, the IPQ-Revised (IPQ-R), which has demonstrated improved psychometric properties [19]. The original IPQ and the IPQ-R have been applied to a broad range of disease and the pre-eminence of the tool within clinical research to evaluate patient illness perceptions has led to the measure being the instrument of choice for clinical research and practice [20,21].

Recognising further the influence of carers, both professional, family members and advocates of the support of patients in coping with disease and engaging with treatment, the IPQ/IPQ-R has been developed into versions that can be readily used in these particular groups [22,23]. The reliability of the instrument applied to these contexts (e.g., clinical staff and carers) has been less extensively evaluated, a surprising observation given the influence of these groups on the patient's response to illness, compliance and engagement with treatment and thus its influence on clinical outcome. Interestingly, within the arena of mental health and major axis I psychiatric disorders, such as schizophrenia, very little research has been conducted to evaluate the measurement properties of the IPQ/IPQ-R, a finding made noteworthy by the recognition of the importance of illness perceptions within recovery models that are highly influential both to the delivery of evidence-informed, person-centered mental healthcare, and to the contextual alignment of treatment intervention to outcome [15,24-27].

## Evaluation of the measurement properties of the IPQ/IPQ-R

The internal consistency of the IPQ has been reported in a number of studies as support for the measurement validity of the scale, measures of Cronbach's coefficient alpha being generally satisfactory, though not always across all of the subscales [5,20,23]. The utility of Cronbach's alpha as a singular index of instrument measurement fidelity is limited both theoretically, for example, (high) internal consistency does not equate to instrument unidimensionality [28-31] and also from a measurement perspective in that Cronbach's alpha is a lower-bound measure of instrument reliability and may significantly underestimate the reliability of a scale [30,32,33].

The structural validity of the IPQ has also been evaluated using principal component analysis, exploratory factor analysis (EFA) and confirmatory factor analysis (CFA), with mixed results. One of the few studies which has evaluated the psychometric properties of the IPQ in clinical staff caring for patients with axis I disorders was

that of Fleming *et al.* [23] using the schizophrenia carers version of the IPQ (IPQ-SCV) developed by Barrowclough and colleagues [22]. Fleming and colleagues [23] found the (IPQ-SCV) to have poor measurement characteristics in this group and suggested the tool was unsuitable for use in this group due to poor concordance of data to anticipated factor structure and low internal consistency as assessed by Cronbach's alpha on a number of IPQ subscales. However, these researchers specifically highlighted these limitations to a specific context of evaluating psychosocial intervention training (the participants in their study) and thus generalising more broadly on the suitability of the tool for use in other clinical groups working with people with schizophrenia may not be appropriate.

The issue of the validity of the IPQ for use in practitioners working with patients with axis I disorders and indeed co-morbid axis II disorder is still, therefore, to be addressed. Recognising this important vacuum in the literature, the current study sought to evaluate the measurement properties of the IPQ-SCV in practitioners working in practice with patients with significant axis I and axis II co-morbidity, specifically schizophrenia co-morbid with learning disability.

## Methods

### Population

Participants (n=210) were a range of mental healthcare workers (both professionally qualified and unqualified) with a mean age of 46.4 years (SD 8.17) and mean length of service of 20.36 years (SD 10.82) working with patients with a dual diagnosis of schizophrenia and learning disability. Fifty-four of the participants worked in community settings, 8 in specialist settings, 102 in in-patient care settings (missing data on 4 participants). The majority of participants were female (n=138). A minority of participants were unqualified (n=71), while the largest group (n=100) were Registered Nurses - Learning Disability (RN-LD). A small number (n=5) of Registered Nurses - Mental Health (RN-MH) and dual-qualified (n=4) nurses (RN-LD & RN-MH) and those with an appropriate vocational qualification or higher national certificate (n=21) comprised the remainder of the participant cohort.

### Ethical approval

Ethical approval and authorisation for the study was provided from the appropriate academic and health organisations prior to commencing data collection.

### Instruments (questionnaires)

The IPQ-SCV [22] was used in the study. This version of the instrument comprised 23 items conceptualized within the original 5-sub-scale model 1: (i) consequences: patient - 7 items; (ii) consequences: relative - 5 items; (iii) control-

cure of illness - 5 items; (iv) control-cure by the practitioner - 2 items & (v) timeline chronic - 2 items, timeline episodic - 2 items.

### Statistical analysis

Replicating the approach taken by Fleming *et al.* [23] 3 measurement models of the IPQ were evaluated using CFA. The first model (model 1) was the original 5-factor model of the IPQ. The second model (model 2) tested split the timeline fact into 2 distinct factors of chronic and episodic as suggested by Barrowclough *et al.* [22], thus presenting a 6-factor model for evaluation. A uni-dimensional model of the IPQ with all items loading on a single factor was also evaluated. A maximum-likelihood (ML) estimation method was adopted with robust standard errors generated using the Satorra-Bentler scaled correction [34]. Consistent with contemporary CFA model evaluation practice, a range of goodness-of-fit tests [35-40] were used to evaluate the specified models.

Internal consistency of each IPQ sub-scale was evaluated using Cronbach's alpha [41] and McDonald's [42,43] omega reliability statistic. The omega hierarchical ( $\omega_h$ ) test statistic [43] has been found to be a more accurate measure of reliability compared to Cronbach's alpha [30]. The omega total ( $\omega_t$ ) test statistic [42] provides an index of

reliability within the context of the total saturation of the test. Hierarchical and total omega levels were calculated based on 5-factor model and a 6-factor model from a minimum residual factor analysis of the dataset followed by a Schmid-Leiman transformation to establish general factor loadings and from which  $\omega_h$  and  $\omega_t$  are calculated. Data were analysed using the R statistical modelling program [44].

### Procedure

Participants are asked how much they agree with the statement in each item on a 5-point scale (Strongly agree to Strongly disagree). Basic demographic questions relating to age, gender, length of service and clinical area were also asked on a supplementary sheet.

### Results

The mean scores on the IPQ subscales are shown in Table 1. Individual item skew and kurtosis characteristics were scrutinised which revealed no evidence of excessive skew (individual item max = 1.95) or kurtosis (individual item max = 3.87).

Table 1 Mean, standard deviation, range and confidence interval (CI) of the Illness Perception Questionnaire sub-scale scores

Variable	Mean	SD	95% CI	Range	Skew	Kurtosis	Alpha
Consequences: relative	11.56	3.03	11.14 - 11.97	14	0	-0.37	0.71
Control: cure of illness	16.7	2.74	16.33 - 17.07	17	-0.42	0.43	0.5
Control: cure by practitioner	7.25	1.5	7.05 - 7.46	8	-0.15	0.52	0.61
Timeline: chronic and episodic	12.45	1.81	12.2 - 12.7	14	-0.23	2.09	0.31
Timeline: chronic	4.04	1.51	3.84 - 4.25	8	0.57	0.51	0.86
Timeline: episodic	8.41	1.44	8.21 - 8.6	8	-1.16	2.58	0.83

Note: Separate chronic and episodic timeline subscales are shown as these are representative of the 6-factor model evaluated by confirmatory factor analysis

Evaluation of the skew and kurtosis characteristics of the scale reveal that the data appear distributionally normal and suitable for an ML estimation approach to CFA.

The factor models evaluated using CFA and accompanying fit indices are shown in Table 2.

Table 2 Evaluation of the structure of the IPQ determined by CFA

Model	$\chi^2$	d.f.	<i>p</i>	CFI	TLI	RMSEA	SRMR
Five-factor	567.08	220	<0.001	0.65	0.6	0.09	0.1
Six-factor	449.67	215	<0.001	0.76	0.72	0.07	0.09
Single factor	881.07	230	<0.001	0.34	0.28	0.12	0.12

Note: Best model fit indices from confirmatory factor analysis indicated in bold IPQ, Illness Perception Questionnaire; CFA, confirmatory factor analysis; RMSEA, root mean squared error of approximation; SRMR, standardised root mean square residual; CFI, comparative fit index; TLI, Tucker-Lewis index

The 6-factor model offered the best fit to the data compared to the 5-factor and single factor models; however, evaluation of the fit indices of all models revealed a consistent pattern of poor model fit. Cronbach's alpha was sub-optimal for a number of the IPQ subscales with the exception of the consequences-relative and

timelines-chronic and timelines-episodic subscales. The Cronbach's alpha of the total scale was also sub-optimal (0.66). Evaluation of the hierarchical and total omega levels are summarised in Table 3. These reveal identical findings for 5- and 6-factor specified models. The  $\omega_t$  statistic, which is relevant to the proportion of variance in

the scale in relation to all common factors was 0.81, whereas the  $\omega_h$  statistic which indicates the variance associated with a common factor, was sub-optimal within a range of 0.32-0.36.

Table 3 Evaluation of omega reliability measures based on the 5- and 6-factor models

Model	$\omega_h$	$\omega_t$
Five-factor	0.32	0.81
Six-factor	0.36	0.81

## Discussion

A combination of schizophrenia with co-morbid learning disability may represent additional challenges for both the patient themselves and staff engaged in optimising patient function within contemporary treatment and recovery models of care. Critical to patient outcome are the attitudes, beliefs and opinions of care staff to both schizophrenia and learning disability. Accurate and reliable assessment of the illness perceptions of care staff working with this group is therefore essential, both for maximising the opportunities for facilitating positive patient outcomes and for identifying potential training and education opportunities for staff in the event of negative or counter-evidence findings of staff illness perception of this group. Central to this activity is an accurate and reliable measure of illness perceptions. However, consistent with the findings of Fleming *et al.* [23] in those professionals working with patients with schizophrenia, the IPQ was found to have unsatisfactory measurement characteristics in the current study, essentially, a poor-fit to contemporary measurement models of the tool and poor overall internal consistency in relation to Cronbach’s alpha. However, the Omega statistics suggest a more complex picture in relation to overall instrument reliability. It was observed that  $\omega_t$  was satisfactory ( $>0.7$ ) for the measure when evaluating the variance due to all common factors, but conversely unsatisfactory ( $\omega_h$ ) when evaluating the variance due to a general factor. This may indicate support for the intrinsic multi-dimensionality of the measure, in contrast to the CFA findings which emphasise that the current conceptual measurement models of the IPQ are miss-specified.

The above findings raise a number of important considerations. Firstly, are the measurement limitations of IPQ specific to this group of care workers working within this clinical population or is this representative of a more generalizable problem with the IPQ? In some respects this question is difficult to address comprehensively, since there are many studies in other groups which emphasise the adequacy of the IPQ. Secondly, the version of the IPQ used in the current study has been adapted for use in a schizophrenia context, therefore representing a modification/adaptation of an existing tool. Consequently, the question can only be addressed in relation to this clinical context (schizophrenia and learning disability) and the participant population. However, comparison with the

findings of the current study with those of Fleming *et al.* [23] reveal a remarkably consistent pattern of results, including similar subscale mean scores and variances, poor fit to data and overall poor internal consistency. The current study thus replicates Fleming *et al.*’s [23] study in terms of design and findings and extends it to a carer group working within the context of a more complex psychopathology (schizophrenia co-morbid with learning disability). However, the salient conclusion regarding the comparison of these two studies is that the IPQ represents a poor index of illness perceptions from a measurement perspective in the context of major axis I disorder. This study consequently yields compelling corroborative evidence for the unsuitability for the use of this tool in this group.

Fleming and colleagues [23] highlighted a vacuum in assessment tool availability in illness perceptions emphasising that there was little alternative to using the IPQ. Giving the findings of the current study, it would seem that the development of a new and contemporary model of illness perceptions, specifically for use within the context of axis I disorders, particularly schizophrenia, is now long overdue. A challenge in the development of such a novel measure would be the lack of availability of a comparison measure or ‘gold standard’.

## Conclusion

The findings from the current study indicate the lack of suitability of the IPQ for such a role. A mixed-method [45] approach may therefore be the most suitable and parsimonious approach, starting from first principles and from both patient and staff qualitative interviews to determine the lived experience of both diagnosis and provision of care. Taking such an approach would reduce contamination due to bias that may accompany preconceptions held regarding illness perceptions in relation to this group while providing a ‘clean sheet’ context to develop the *a priori* specified domains and individual items that would comprise the new scale. Taking such an approach, rooted in the evidence-base and derived from the accounts of experts by experience and experts by care provision, would be an appropriate and robust approach toward the development of a disease-specific measure of illness perceptions that could be pursued with methodological robustness, psychometric confidence and clinical appropriateness.

## Conflicts of Interest

The authors declare no conflicts of interest.

## References

[1] Anagnostopoulos, F. & Spanea, E. (2005). Assessing illness representations of breast cancer: a comparison of

- patients with healthy and benign controls. *Journal of Psychosomatic Research* 58 (4) 327-334.
- [2] Henderson, M., Safa, F., Easterbrook, P. & Hotopf, M. (2005). Fatigue among HIV-infected patients in the era of highly active antiretroviral therapy. *HIV Medicine* 6 (5) 347-352.
- [3] Petrie, K.J., Cameron, L.D., Ellis, C.J., Buick, D. & Weinman, J. (2002). Changing illness perceptions after myocardial infarction: an early intervention randomized controlled trial. *Psychosomatic Medicine* 64 (4) 580-586.
- [4] Petrie, K.J., Weinman, J., Sharpe, N. & Buckley, J. (1996). Role of patients' view of their illness in predicting return to work and functioning after myocardial infarction: longitudinal study. *British Medical Journal* 312 (7040) 1191-1194.
- [5] Weinman, J., Petrie, K.J., Moss-Morris, R. & Horne, R. (1996). The Illness Perception Questionnaire: a new method for assessing the cognitive representations of illness. *Psychology & Health* 11 (3) 431-445.
- [6] Loignon, C., Bedos, C., Sevigny, R. & Leduc, N. (2009). Understanding the self-care strategies of patients with asthma. *Patient Education and Counseling* 75 (2) 256-262.
- [7] Millar, K., Purushotham, A.D., McLatchie, E., George, W.D. & Murray, G.D. (2005). A 1-year prospective study of individual variation in distress, and illness perceptions, after treatment for breast cancer. *Journal of Psychosomatic Research* 58 (4) 335-342.
- [8] Werner, S. (2012). Intention to work with individuals with dual diagnosis: testing the theory of planned behavior among students from various professions. *Health & Social Work* 37 (2) 81-89.
- [9] Wray, J., Lindsay, B., Crozier, K., Andrews, L. & Leeson, J. (2013). Exploring perceptions of psychological services in a children's hospice in the United Kingdom. *Palliative & Supportive Care* 11 (5) 373-382.
- [10] Chan, J.C., Ong, J.C., Avalos, G., Regan, P.J., McCann, J., Groarke, A. & Kelly, J.L. (2009). Illness representations in patients with hand injury. *Journal of Plastic Reconstructive & Aesthetic Surgery* 62 (7) 927-932.
- [11] French, D.P., Wade, A.N., Yudkin, P., Neil, H.A., Kinmonth, A.L. & Farmer, A.J. (2008). Self-monitoring of blood glucose changed non-insulin-treated Type 2 diabetes patients' beliefs about diabetes and self-monitoring in a randomized trial. *Diabetic Medicine* 25 (10) 1218-1228.
- [12] Lee, B.O., Chien, C.S., Hung, C.C. & Chou, P.L. (2015). Effects of an in-hospital nursing intervention on changing illness perceptions in patients with injury. *Journal of Advanced Nursing* doi: 10.1111/jan.12716.
- [13] Quiles Marcos, Y., Terol Cantero, M.C., Romero Escobar, C. & Pagan Acosta, G. (2007). Illness perception in eating disorders and psychosocial adaptation. *European Eating Disorders Review* 15 (5) 373-384.
- [14] Cook, S.A., Salmon, P., Dunn, G., Holcombe, C., Cornford, P. & Fisher, P. (2015). The association of metacognitive beliefs with emotional distress after diagnosis of cancer. *Health Psychology* 34 (3) 207-215.
- [15] Higbed, L. & Fox, J.R. (2010). Illness perceptions in anorexia nervosa: a qualitative investigation. *British Journal of Clinical Psychology* 49 (Part 3) 307-325.
- [16] Israel, J.I., White, K.S. & Gervino, E.V. (2015). Illness perceptions, negative emotions, and pain in patients with noncardiac chest pain. *Journal of Clinical Psychology in Medical Settings* 22 (1) 77-89.
- [17] Kaptein, A.A., Scharloo, M., Fischer, M.J., Snoei, L., Cameron, L.D., Sont, J.K., Rabe, K.F. & Weinman, J. (2008). Illness perceptions and COPD: an emerging field for COPD patient management. *Journal of Asthma* 45 (8) 625-629.
- [18] Katerndahl, D.A. & Parchman, M. (2002). The ability of the stress process model to explain mental health outcomes. *Comprehensive Psychiatry* 43 (5) 351-360.
- [19] Moss-Morris, R., Weinman, J., Petrie, K.J., Horne, R., Cameron, L.D. & Buick, D. (2002). The Revised Illness Perception Questionnaire (IPQ-R). *Psychology & Health* 17, 1-16.
- [20] Leysen, M., Nijs, J., Meeus, M., Paul van Wilgen, C., Struyf, F., Vermandel, A., Kuppens, K. & Roussel, N.A. (2015). Clinimetric properties of illness perception questionnaire revised (IPQ-R) and brief illness perception questionnaire (Brief IPQ) in patients with musculoskeletal disorders: A systematic review. *Manual Therapy* 20 (1) 10-17.
- [21] Maas, M., Taal, E., van der Linden, S. & Boonen, A. (2009). A review of instruments to assess illness representations in patients with rheumatic diseases. *Annals of Rheumatic Diseases* 68 (3) 305-309.
- [22] Barrowclough, C., Lobban, F., Hatton, C. & Quinn, J. (2001). An investigation of models of illness in carers of schizophrenia patients using the Illness Perception Questionnaire. *British Journal of Clinical Psychology* 40 (Part 4) 371-385.
- [23] Fleming, M.P., Martin, C.R., Miles, J. & Atkinson, J. (2009). The utility of the Illness Perception Questionnaire in the evaluation of mental health practitioners' perspectives on patients with schizophrenia. *Journal of Evaluation in Clinical Practice* 15 (5) 826-831.
- [24] Barrett, B., Young, M.S., Teague, G.B., Winarski, J.T., Moore, K.A. & Ochshorn, E. (2010). Recovery orientation of treatment, consumer empowerment, and satisfaction with services: a mediational model. *Psychiatric Rehabilitation Journal* 34 (2) 153-156.
- [25] Romano, D.M., McCay, E., Goering, P., Boydell, K. & Zipursky, R. (2010). Reshaping an enduring sense of self: the process of recovery from a first episode of schizophrenia. *Early Intervention in Psychiatry* 4 (3) 243-250.
- [26] Russinova, Z., Rogers, E.S., Ellison, M.L. & Lyass, A. (2011). Recovery-promoting professional competencies: perspectives of mental health consumers, consumer-providers and providers. *Psychiatric Rehabilitation Journal* 34 (3) 177-185.
- [27] Werner, S. (2012). Subjective well-being, hope, and needs of individuals with serious mental illness. *Psychiatry Research* 196 (2-3) 214-219.
- [28] Green, S.B., Lissitz, R.W. & Mulaik, S.A. (1977). Limitations of coefficient alpha as an index of test unidimensionality. *Educational and Psychological Measurement* 37, 827-838.
- [29] Nunnally, J.C. (1978). *Psychometric Theory*. 2nd edn. New York: McGraw-Hill.

- [30] Zinbarg, R.E., Revelle, W., Yovel, I. & Li, W. (2005). Cronbach's  $\alpha$ , Revelle's  $\beta$ , and McDonald's  $\omega_h$  : their relations with each other and two alternative conceptualizations of reliability. *Psychometrika* 70, 123-133.
- [31] Sijtsma, K. (2009). On the Use, the Misuse, and the Very Limited Usefulness of Cronbach's Alpha. *Psychometrika* 74 (1) 107-120.
- [32] Eisinga, R., te Grotenhuis, M. & Pelzer, B. (2013). The reliability of a two-item scale: Pearson, Cronbach or Spearman-Brown? *International Journal of Public Health* 58, 637-642.
- [33] Graham, J.M. (2006). Congeneric and (essentially) tau-equivalent estimates of score reliability: What they are and how to use them. *Educational and Psychological Measures* 66, 930-944.
- [34] Satorra, A. & Bentler, P.M. (2001). A scaled difference chi-square test statistic for moment structure analysis. *Psychometrika* 66, 507-514.
- [35] Bentler, P.M. (1990). Comparative fit indexes in structural models. *Psychological Bulletin* 107 (2) 238-246.
- [36] Bentler, P.M. & Bonett, D.G. (1980). Significance tests and goodness of fit in the evaluation of covariance structures. *Psychological Bulletin* 88, 588-606.
- [37] Hu, L.T. & Bentler, P.M. (1995). Evaluating model fit. In: Structural Equation Modelling: Concepts, Issues and Applications. Hoyle, R.H., ed. Thousand Oaks, CA: Sage.
- [38] Kline, R.B. (1998). Principles and Practice of Structural Equation Modeling. New York: Guilford Press.
- [39] Marsh, H.W., Balla, J.R. & McDonald, R.P. (1998). Goodness-of-fit indices in confirmatory factor analysis: the effect of sample size. *Psychological Bulletin* 103, 391-410.
- [40] Tucker, L.R. & Lewis, C. (1973). A reliability coefficient for maximum likelihood factor analysis. *Psychometrika* 38, 1-10.
- [41] Cronbach, L.J. (1951). Coefficient alpha and the internal structure of tests. *Psychometrika* 16, 297-334.
- [42] McDonald, R.P. (1978). Generalizability in factorable domains: "Domain validity and generalizability". *Educational and Psychological Measures* 38, 75-79.
- [43] McDonald, R.P. (1999). Test Theory: A Unified Treatment. Mahwah, NJ: Erlbaum.
- [44] Team, R.C. (2013). R: A Language and Environment for Statistical Computing. Vienna, Austria: Foundation for Statistical Computing.
- [45] Creswell, J.W. & Plano Clark, V.L. (2011). Designing and Conducting Mixed Methods Research. 2nd edn. Thousand Oaks, CA: Sage.